

CS5371

Theory of Computation

Lecture 1: Mathematics Review I
(Basic Terminology)

Objectives

- Unlike other CS courses, this course is a MATH course...
- We will look at a lot of definitions, theorems and proofs
- This lecture: reviews basic math notation and terminology
 - Set, Sequence, Function, Graph, String...
- Also, common proof techniques
 - By construction, induction, contradiction

Set

- A **set** is a group of items
- One way to describe a set: list every item in the group inside { }
 - E.g., { 12, 24, 5 } is a set with three items
- When the items in the set has trend: use ...
 - E.g., { 1, 2, 3, 4, ... } means the set of natural numbers
- Or, state the rule
 - E.g., { $n \mid n = m^2$ for some positive integer m } means the set { 1, 4, 9, 16, 25, ... }
- A set with no items is an **empty set** denoted by {} or \emptyset

Set

- The order of describing a set does not matter
 - $\{12, 24, 5\} = \{5, 24, 12\}$
- Repetition of items does not matter too
 - $\{5, 5, 5, 1\} = \{1, 5\}$
- Membership symbol \in
 - $5 \in \{12, 24, 5\}$ $7 \notin \{12, 24, 5\}$

Set (Quick Quiz)

- How many items are in each of the following set?
 - $\{ 3, 4, 5, \dots, 10 \}$
 - $\{ 2, 3, 3, 4, 4, 2, 1 \}$
 - $\{ 2, \{2\}, \{\{2\}\} \}$
 - \emptyset
 - $\{\emptyset\}$

Set

Given two sets A and B

- we say $A \subseteq B$ (read as A is a **subset** of B) if every item in A also appears in B
 - E.g., A = the set of primes, B = the set of integers
- we say $A \subsetneq B$ (read as A is a **proper subset** of B) if $A \subseteq B$ but $A \neq B$

Warning: Don't be confused with \in and \subseteq

- Let $A = \{1, 2, 3\}$. Is $\emptyset \in A$? Is $\emptyset \subseteq A$?

Union, Intersection, Complement

Given two sets A and B

- $A \cup B$ (read as the **union** of A and B) is the set obtained by combining all elements of A and B in a single set
 - E.g., $A = \{1, 2, 4\}$ $B = \{2, 5\}$
 $A \cup B = \{1, 2, 4, 5\}$
- $A \cap B$ (read as the **intersection** of A and B) is the set of common items of A and B
 - In the above example, $A \cap B = \{2\}$
- \bar{A} (read as the **complement** of A) is the set of items under consideration not in A

Set

- The **power set** of A is the set of all subsets of A , denoted by 2^A
 - E.g., $A = \{0, 1\}$
$$2^A = \{ \{\}, \{0\}, \{1\}, \{0,1\} \}$$
 - How many items in the above power set of A ?
- If A has n items, how many items does its power set contain? Why?

Sequence

- A **sequence** of items is a list of these items in some order
- One way to describe a sequence: list the items inside ()
 - (5, 12, 24)
- Order of items inside () matters
 - (5, 12, 24) \neq (12, 5, 24)
- Repetition also matters
 - (5, 12, 24) \neq (5, 12, 12, 24)
- Finite sequences are also called **tuples**
 - (5, 12, 24) is a 3-tuple
 - (5, 12, 12, 24) is a 4-tuple

Sequence

Given two sets A and B

- The **Cartesian product** of A and B , denoted by $A \times B$, is the set of all possible 2-tuples with the first item from A and the second item from B
 - E.g., $A = \{1, 2\}$ and $B = \{x, y, z\}$
 $A \times B = \{ (1,x), (1,y), (1,z), (2,x), (2,y), (2,z) \}$
- The Cartesian product of k sets, A_1, A_2, \dots, A_k , denoted by $A_1 \times A_2 \times \dots \times A_k$, is the set of all possible k -tuples with the i^{th} item from A_i

Functions

- A **function** takes an input and produces an output
- If f is a function, which gives an output b when input is a , we write
$$f(a) = b$$
- For a particular function f , the set of all possible input is called f 's **domain**
- The outputs of a function come from a set called f 's **range**

Functions

- To describe the property of a function that it has domain D and range R , we write

$$f : D \rightarrow R$$


- E.g., The function `add` (to add two numbers) will have an input of two integers, and output of an integer
 - We write: `add: $\mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}$`

Functions (Quick Quiz)

- *Guess:* What does the following function DOW do?
 - $DOW(9,12) = 2$
 - $DOW(9,13) = 3$
 - $DOW(9,17) = 7$
 - $DOW(9,18) = 1$
- What are the domain and the range of DOW?

Graphs

- A graph is a set of points with lines connecting some of the points
- Points are called **vertices**, lines are called **edges**
- E.g.,




Graphs

- The number of edges at a particular vertex is the **degree** of the vertex
- In the previous example, 3 vertices have degree = 2
- A graph can be described by telling what are its vertices, and what are its edges. Formally, a graph G can be written as $G = (V, E)$, where V is the set of vertices, and E is the set of edges

Graphs

- We say a graph G is a **subgraph** of H if vertices of G are a subset of the vertices of H , and all edges in G are the edges of H on the corresponding vertices



Graph H


Subgraph G
shown darker

Graphs

- A **path** is a sequence of vertices connected by edges
- If every two nodes have a path between them, the graph is **connected**
- A **cycle** is a path that starts and ends at the same vertex
- A **tree** is a connected graph with no cycles

Graphs (Quick Quiz)

- Is the following graph connected?
- Is it a tree?
- Are there any cycles?
- How about the darker subgraph?




Directed Graphs

- If lines are replaced by arrows, the graph becomes **directed**
- The number of arrows pointing into a vertex is called **in-degree** of the vertex
- The number of arrows pointing from a vertex is called **out-degree** of the vertex
- A **directed path** is a path from one vertex to the other vertex, following the direction of the "arrows"

Directed Graphs

- Is there a directed path from a to b?



Strings

- An **alphabet** = a set of characters
 - E.g., The English Alphabet = {A,B,C,...,Z}
- A **string** = a sequence of characters
- A string *over* an alphabet Σ
 - A sequence of characters, with each character coming from Σ
- The **length** of a string w , denoted by $|w|$, is the number of characters in w
- The **empty string** (written as ε) is a string of length 0

Strings

Let $w = w_1w_2\dots w_n$ be a string of length n

- A **substring** of w is a consecutive subsequence of w (that is, $w_iw_{i+1}\dots w_j$ for some $i \leq j$)
- The **reverse** of w , denoted by w^R , is the string $w_n\dots w_2 w_1$
- A set of strings is called a **language**

Next time

- Common Proof Techniques
- Part I: Automata Theory